



Review

Suppression of soil nitrification by plants

Guntur Venkata Subbarao^{a,*}, Tadashi Yoshihashi^a, Margaret Worthington^b,
Kazuhiko Nakahara^a, Yasuo Ando^a, Kanwar Lal Sahrawat^c,
Idupulapati Madhusudhana Rao^b, Jean-Christophe Lata^d, Masahiro Kishii^e,
Hans-Joachim Braun^e

^a Japan International Research Center for Agricultural Sciences (JIRCAS), 1-1 Ohwashi, Tsukuba, Ibaraki 305-8686, Japan

^b Centro Internacional de Agricultura Tropical (CIAT), A.A. 6713, Cali, Colombia

^c International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Hyderabad, Andhra Pradesh, India

^d Sorbonne Universities, UPMC Univ. Paris 06, UMR 7618, Institut d'Économie Supérieure, 46 rue d'Ulm, 75230 Paris Cedex, France;

^e Department of Geocology and Geochemistry, Institute of Natural Resources, Tomsk Polytechnic University, 30, Lenin Street, Tomsk, 634050, Russia

^e CIMMYT (International Maize and Wheat Improvement Center), Apdo Postal 6-641, 06600 Mexico, D.F., Mexico

ARTICLE INFO

Article history:

Received 21 October 2014

Received in revised form

22 December 2014

Accepted 21 January 2015

Available online 28 January 2015

Keywords:

Biological nitrification inhibition (BNI)

Climate change

Global warming

Nitrification inhibitors

Nitrous oxide emissions

Nitrogen use efficiency (NUE)

ABSTRACT

Nitrification, the biological oxidation of ammonium to nitrate, weakens the soil's ability to retain N and facilitates N-losses from production agriculture through nitrate-leaching and denitrification. This process has a profound influence on what form of mineral-N is absorbed, used by plants, and retained in the soil, or lost to the environment, which in turn affects N-cycling, N-use efficiency (NUE) and ecosystem health and services. As reactive-N is often the most limiting in natural ecosystems, plants have acquired a range of mechanisms that suppress soil-nitrifier activity to limit N-losses via N-leaching and denitrification. Plants' ability to produce and release nitrification inhibitors from roots and suppress soil-nitrifier activity is termed 'biological nitrification inhibition' (BNI). With recent developments in methodology for *in-situ* measurement of nitrification inhibition, it is now possible to characterize BNI function in plants. This review assesses the current status of our understanding of the production and release of biological nitrification inhibitors (BNIs) and their potential in improving NUE in agriculture. A suite of genetic, soil and environmental factors regulate BNI activity in plants. BNI-function can be genetically exploited to improve the BNI-capacity of major food- and feed-crops to develop next-generation production systems with reduced nitrification and N₂O emission rates to benefit both agriculture and the environment. The feasibility of such an approach is discussed based on the progresses made.

© 2015 Elsevier Ireland Ltd. All rights reserved.

Contents

| | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 1. Introduction | 156 |
| 1.1. Nitrification: A biological process of critical importance for the sustainability of agricultural systems with implications for climate change | 156 |
| 2. Biological nitrification inhibition (BNI) | 156 |
| 2.1. The BNI concept | 156 |
| 2.2. Evidence for BNI-function | 157 |
| 2.3. Chemical identity of BNIs and their mode of action | 158 |
| 2.4. Characterization of BNI-function | 158 |
| 2.5. BNI release mechanisms | 159 |
| 2.6. Stability of BNIs in soil systems | 160 |
| 2.7. Potential for genetic improvement of BNI-capacity | 161 |
| 2.8. Deploying BNI function in production agriculture | 161 |

* Corresponding author. Tel.: +81 29 838 6354; fax: +81 298386354.

E-mail addresses: subbarao@jircas.affrc.go.jp, subbagv2000@gmail.com (G.V. Subbarao).




Fig. 1. Major processes of nitrogen cycle in soils.

H⁺ is released by plant roots when NH₄⁺ is absorbed and assimilated, leading to the acidification of the rhizosphere, which improves P availability to growing plants [45].

Agronomic NUE (NUE_{agronomic} = grain yield per unit of applied N) is a function of both intrinsic-NUE (NUE_{intrinsic} = dry matter produced per unit N absorbed), HI (harvest index) and N uptake [46]. NUE_{intrinsic} is physiologically conserved [47]; thus, improvement in NUE_{agronomic} can only come from improvements in crop-N uptake, which is largely a function of recovering applied fertilizer-N [48]. BNI function in plants impacts NUE_{agronomic} by improving N uptake, facilitate N-retention and reduce N-losses associated with nitrification–denitrification processes [30,49], a conclusion further supported by both field and modeling studies [44,50–53].

Recent methodological developments have facilitated the detection and quantification of nitrification inhibitors from plant roots using a recombinant luminescent *Nitrosomonas* construct [54,55]. The recombinant strain of *N. europaea* carries an expression vector for the *Vibrio harveyi luxAB* genes (Fig. 2) and produces a distinct two-peak luminescence pattern during a 30-s analysis period [55]. The functional relationship between bioluminescence emission and nitrite production in the assay has been shown to be linear using a synthetic nitrification inhibitor, allylthiourea (AT) [55]. The inhibition caused by 0.22 μM AT in assay (about 80% inhibition in bioluminescence and NO₂⁻ production) is defined as one allylthiourea unit (ATU) [55]. Using the response to a concentration gradient of AT (i.e., standard dose-response curve), the inhibitory effects of root exudates, soil or plant extracts are determined and expressed in ATU [55]. These research methodologies have facilitated the evaluation and characterization of BNI-function in plants [55]. Soil-based assays to determine changes in nitrification potential in the rhizosphere [56] and analysis of nitrifier populations

can further complement these efforts to characterize BNI function [49,57].

2.2. Evidence for BNI-function

Most plants release chemical compounds from root systems that either stimulate or suppress nitrifier activity. The assay system based on recombinant luminescent *Nitrosomonas* can be used to detect and quantify nitrification inhibitors (i.e. BNI-activity) or stimulators (i.e. negative BNI-activity) released from roots [54,55]. The root exudates of most legumes (*Glycine max*, *Vigna unguiculata* and *Phaseolus vulgaris*) did not have detectable inhibitory activity in the assay, whereas most cereals evaluated have varying levels of inhibitory activity in the assay [58]. Isoflavones such as genistein and daidzein found in soybean root exudates have stimulatory effect on *Nitrosomonas* when tested in the assay (GV Subbarao, unpublished results). Tropical pasture grasses that are adapted to low-N environments, in particular *Brachiaria* spp. have the highest BNI-activity in root systems [58]. In contrast, *Panicum* spp. adapted to high-N (in comparison to *Brachiaria* spp.) environments have relatively weak BNI-activity in their root systems [58,59]. Among field crops, sorghum (*Sorghum bicolor*) adapted to low-N input environments appears to have stronger BNI-capacity than crops adapted to high-N input environments such as wheat (*Triticum aestivum*) and maize (*Zea mays*) [58]. Nitrification inhibition is likely an adaptation mechanism to retain and use N efficiently in N limiting natural systems [30,52,59,60]. It is not surprising that N-fixing legumes have low BNI-capacity in root systems as BNI-function may have no adaptive value to them; indeed BNI may favor the attraction of non-legume competitors in N-limiting environments [58]. However, the intensity of N-fixation itself depends on N availability [61], and therefore should be modified by the BNI-capacity of other




Fig. 2. Map of recombinant luminous *Nitrosomonas europaea* (pHUX20) developed to detect and quantify nitrification inhibitors in the plant–soil system [54].

plants, which raises questions about the evolution and plasticity of this process in complex plant communities.

2.3. Chemical identity of BNIs and their mode of action

Several BNIs have been isolated from root exudates and plant tissues (root and shoot) of *B. humicola* and sorghum [49,55,62–64] (Table 1). These BNIs have chemical structures belonging to diverse functional groups – fatty acids [linoleic acid, and linolenic acid], phenylpropanoids [Methyl 3-(4-hydroxyphenyl)propionate (MHPP), methyl-*p*-coumarate, and methyl ferulate], flavonoids [sakuranetin and karanjin, quinones [sorgoleone], diterpenoids [brachialactone] [30] and isothiocyanates [2-propenyl-glucosinolate, methyl-isothiocyanate, 2-propenyl-isothiocyanate, butyl-isothiocyanate, phenyl-isothiocyanate, benzyl-isothiocyanate, butyl-isothiocyanate, phenyl-isothiocyanate, benzyl-isothiocyanate and phenethyl-isothiocyanate] [68–70].

Dominant compounds such as hydrophilic-brachialactone released from *B. humicola* roots or hydrophobic-sorgoleone released from sorghum roots account for a major portion (>80%) of BNI-activity in those species [30,49,65]. The sorgoleone biosynthetic pathway is known and its genetic control is well-understood [71,72]. In contrast, the biosynthesis pathway for brachialactone is still unknown. Brachialactone has a dicyclopenta[a,d]cyclooctane skeleton (5–8–5 ring system) with a γ -lactone ring bridging one of the five-membered rings and the eight membered ring [49,73]. Certain fungi and plants have the ability to synthesize 5–8–5 tricyclic terpenoids such as ophiobolanes and fusicoccanes [73–75], but the lactone ring of brachialactone is a novel cyclic diterpenoid. Fusicoccin type cyclic diterpenes are biologically synthesized from geranylgeranyl diphosphate by a two-step cyclization catalyzed by terpene cyclases [75]. In higher plants, terpenoid biosynthesis is through either HMG-CoA reductase pathway (mevalonic acid pathway, located in cytoplasm) or 2-C-methyl-D-erythritol

4-phosphate/1-deoxy-D-xylulose 5-phosphate pathway (MEP/DOXP pathway, located in plastids) [76,77]. Operon-like gene clusters control the biosynthesis pathways of certain diterpenoids such as ‘momilactone’ in rice [76–80], or certain phytoalexins such as sakuranetin [81]. If such an operon cluster for brachialactone biosynthesis is identified, metabolic engineering of brachialactone biosynthesis and introduction of BNI-capacity into root systems of major food crops using transgenic approaches may be possible.


Crude BNI-activity extracted from root exudates or plant tissues is likely composed of a cocktail of nitrification inhibitors, each with a single mode or in some cases multi-mode of inhibitory effects on enzymatic pathways of *Nitrosomonas* [30]. BNIs such as linoleic acid, linolenic acid, sorgoleone and brachialactone inhibit *Nitrosomonas* through blocking of both ammonia mono-oxygenase and hydroxylamine oxidoreductase enzymatic pathways involved in ammonia oxidation in *Nitrosomonas* [49,62,65]. Also, some BNIs (e.g. sorgoleone) could disrupt the crucial electron transfer pathway from hydroxylamine oxidoreductase to ubiquinone and cytochrome. This pathway needs to be maintained to generate reducing power (i.e. NADPH), which is crucial to the metabolic functions of *Nitrosomonas* [82–84]. Most synthetic nitrification inhibitors [(e.g., nitrapyrin, dicyandiamide, and 3,4-dimethylpyrazole phosphate (DMPP)] inhibit *Nitrosomonas* activity by suppressing the ammonia monooxygenase enzymatic pathway [60,85], but they have no effect on the hydroxylamine oxidoreductase enzymatic pathway. BNIs such as hydrophilic-MHPP and certain monoterpenes (e.g. limonene, produced by *Pinus ponderosa*) also inhibit *Nitrosomonas* in a similar way, i.e. by blocking only the ammonia monooxygenase pathway [22,63,67].

2.4. Characterization of BNI-function

Two categories of BNIs are released from plant root systems: hydrophilic-BNIs and hydrophobic-BNIs (Fig. 3) [65]. Their

Table 1Relative effectiveness of various BNIs, and their mode of action on *Nitrosomonas* in *in vitro* bioassay (AMO, ammonia monooxygenase; HAO, hydroxyl aminooxidoreductase).

| Serial No. | BNI compound | Isolated from | Inhibit AMO or HAO enzymatic pathway | ED ₈₀ [(μM) in <i>in vitro</i> bioassay] | Ref. |
|------------------------------------|----------------------------|-----------------------------------|--------------------------------------|-----------------------------------------------------|---------|
| 1 | Brachialactone | <i>B. humidicola</i> root exudate | AMO and HAO | 10.6 | [49] |
| 2 | Methyl <i>p</i> -coumarate | <i>B. humidicola</i> root tissue | NA | >40.0 | [64] |
| 3 | Methyl ferulate | <i>B. humidicola</i> root tissue | NA | >20.0 | [64] |
| 4 | Linoleic acid (LA) | <i>B. humidicola</i> shoot tissue | AMO and HAO | 16.0 | [62] |
| 5 | Linolenic acid (LN) | <i>B. humidicola</i> shoot tissue | AMO and HAO | 16.0 | [62] |
| 6 | Sorgoleone | Sorghum root exudate | AMO and HAO | 12.0 | [65,66] |
| 7 | MHPP | Sorghum root exudate | AMO | >120.0 | [63] |
| 8 | Sakuranetin | Sorghum root exudate | AMO and HAO | 0.6 | [65] |
| 9 | Limonene | <i>Pinus ponderosa</i> leaf | AMO | NA | [67] |
| Synthetic nitrification inhibitors | | | | | |
| 10 | ©Allylthiourea | | AMO | 0.22 | [62] |
| 11 | ©Nitrapyrin | | AMO | 17.32 | [62] |
| 12 | ©Dicyandiamide | | AMO | 2200.00 | [62] |

**Fig. 3.** Hydrophobic- and hydrophilic-nitrification inhibitors (BNIs) released from plant roots and their significance to BNI function.

relative contribution to BNI-capacity varies among plant species and even across growth stages. For example, in sorghum hydrophobic-BNI activity is the major contributor during early growth stages [up to 14 days after planting], but hydrophobic- and hydrophilic-BNIs contribute equally at later growth stages (30 days and after) [30,65]. Hydrophobic- and hydrophilic-BNIs differ in their mobility in soil due to differential solubility and/or affinity to water. The hydrophobic-BNIs may remain close to the root systems as they are strongly sorbed to soil mineral or organic particles, which may further increase their persistence. Their movement in soil is primarily *via* diffusion across concentration gradients and is likely confined to the rhizosphere [65,82]. In contrast, water soluble hydrophilic-BNIs are more likely to move out of the rhizosphere which may enhance their capacity to suppress nitrification in bulk soil [65]. Thus the distribution of hydrophobic- and hydrophilic-BNIs in the rhizosphere may have complementary functional roles [65].

Based on the observations of BNI-activity in sorghum in several greenhouse studies, it is estimated that the total amount of

BNIs (hydrophilic- plus hydrophobic-) released during 130-days growing period (coinciding with the physiological maturity of this crop) can suppress nitrification up to 50% in about 500 g soil per plant [65]. Field and greenhouse studies with *Brachiaria humidicola* provide evidence for strong nitrification inhibition potential. Assuming 1.5 Mg ha⁻¹ average live root biomass from a long-term grass pasture [86], and BNI-capacity of 17–70 ATU g⁻¹ root dry wt. d⁻¹ [59], it is estimated that *B. humidicola* can potentially release 2.6 × 10⁶ to 7.5 × 10⁶ ATU ha⁻¹ d⁻¹ of hydrophilic-BNIs 49,59; no published results are available on the estimates of hydrophobic-BNI activity from *Brachiaria* spp.). This estimate amounts to an inhibitory potential equivalent to that by the application of 6.2–18 kg of nitrapyrin ha⁻¹ y⁻¹ (based on 1 ATU being equivalent to 0.6 μg of nitrapyrin, a synthetic nitrification inhibitor), which is sufficient to have a significant influence on nitrifier activity and nitrification rates in the soil [49]. Field studies indicate a 90% decline in soil ammonium oxidation rates and N₂O emissions within three years of establishment of *B. humidicola* pastures [49]. These reduced emissions are attributed to the extremely small nitrifier populations present in the established pastures. In the same study, field plots planted to soybean (*Glycine max*, a plant species with no significant BNI-capacity) did not inhibit soil ammonium oxidation rates or N₂O emissions [49] (Fig. 4A and B). Based on the monitoring of N₂O emissions over a 3-year period from fields planted with tropical grasses with a wide range of BNI-capacity, a negative relationship was observed between the BNI-capacity of a species and N₂O emissions [30].

2.5. BNI release mechanisms

BNI synthesis and release are highly regulated plant attributes, which are stimulated by the presence of NH₄⁺ in the rhizosphere [59]. The N-form (NH₄⁺ vs. NO₃⁻) in the soil has a major influence on the synthesis and release of BNIs by *B. humidicola*, sorghum, and *Leymus racemosus*, a wild relative of wheat [49,59,63,65,87,88]. Plants grown with NO₃⁻ as their N-source did not release BNIs, whereas plants grown with NH₄⁺ as the N-source did release BNIs. Despite high levels of BNIs detected in the root tissues of NH₄⁺ grown plants, BNIs were only released when plant roots were directly exposed to NH₄⁺ during the collection of root exudates [49,59,65]. In addition, BNIs release from roots is a localized phenomenon confined to the part of the root system exposed to NH₄⁺ and was not extended to the remaining parts of the root system (Fig. 5) [49,88]. A localized release of BNIs ensures relatively high concentrations of BNIs in the soil micro-sites where nitrifiers are active and NH₄⁺ is present [49]. The regulatory role of NH₄⁺ in the synthesis and release of BNIs suggests a possible adaptive role in protecting NH₄⁺ from nitrifiers [49].

The activation and operation of proton pumping activity of root plasma membranes has been hypothesized as a functional




Fig. 4. (A) Soil ammonium oxidation rates ($\text{mg of NO}_2\text{-N kg}^{-1}\text{ soil d}^{-1}$) in field plots planted with tropical pasture grasses (differing in BNI capacity) and soybean (lacking BNI capacity in roots) [over 3 years from establishment of pastures (September 2004 to November 2007); for soybean, two planting seasons every year and after six seasons of cultivation]. CON, control (plant-free) plots; SOY, soybean; PM, *P. maximum*; BHM, *Brachiaria* hybrid cv. Mulato; BH-679, *B. humidicola* CIAT 679 (standard cultivar); BH-16888, *B. humidicola* accession CIAT 16888 (a germplasm accession). Values are means \pm SE from three replications [49]. (B) Cumulative N_2O emissions ($\text{mg of N}_2\text{O-N m}^{-2}\text{ per year}$) from field plots of tropical pasture grasses (monitored monthly over a 3-year period, from September 2004 to November 2007). Plots are identified in Fig. 3 legend. Values of means \pm SE from three replications [49].

link between BNI release (presumably organic anions) and NH_4^+ uptake and assimilation. If BNIs are transported through voltage-dependent anion channels, their release will be closely related to the regulation of proton pump-ATPase. We speculate that the transport of BNIs, driven by proton pump-ATPase, is associated with NH_4^+ uptake and assimilation in sorghum (Fig. 6) [88]. The rhizosphere pH also influences the release of BNIs from roots. Recent results indicate that sorghum plants do not release BNIs from their roots in the presence of NH_4^+ when the rhizosphere pH is 7 or higher; the optimum BNI release was observed at a rhizosphere pH of 5.0–6.0, which stimulates the functioning of the proton pumps [63,65]. These results imply that the suppression of nitrification by BNI is likely to be restricted to sorghum grown on acid soils. Light-textured soils with low buffering capacity and moderate acidity (pH <6.0), which is the case for most tropical grasslands or savannas, might be better suited for the expression and exploitation of BNI function in sorghum [65,89]. The results of recent studies suggest that unlike hydrophilic-BNI release in sorghum, hydrophobic-BNI release is not sensitive to pH changes in the rhizosphere as its release is not associated with proton pumping activity (Tingjun and Subbarao, unpublished results). Moreover, nitrifier activity and nitrification are suppressed by *B. humidicola* pasture in heavy black




Fig. 5. A hypothesis proposed for localized release of BNIs from roots when NH_4^+ is sensed in the rhizosphere [49].




Fig. 6. A hypothesis on the transport of BNIs, driven by PM H^+ -ATPase, associated with NH_4^+ uptake and assimilation in source [88].

soils (Vertisols) with pH of 7.2 following two years after its establishment [49].

2.6. Stability of BNIs in soil systems

Although quantification of BNI-activity using an *in vitro* assay during 30 min exposure to pure cultures of *Nitrosomonas* spp. is a useful initial screening tool for determining BNI-capacity, we should expect that not all BNIs released from root system of plants are effective in suppressing soil nitrifier activity in the field. For BNIs to be effective in soil-based systems, these compounds must persist and be effective in the soil. BNI-compounds isolated from root exudates of *B. humidicola* were effective when added to the soil at $10\text{--}20\text{ ATU g}^{-1}$ soil and incubated for 55 days at 20°C ; soil nitrification was suppressed by 50–90% [55,90]. Certain BNIs such as linoleic acid and linolenic acid (BNI-compounds isolated from leaf tissues of *B. humidicola*) partially lost their effectiveness in soil after

nitrification/denitrification in soils should be tightly controlled and regulated) to reduce the need for continued application of industrially fixed-N to support food production.

Accelerated nitrification rates in agricultural soils have resulted in a decline in NUE since the advent of Green Revolution, and led to diminishing returns on N-fertilization [32,98,104]. Nearly 70% of applied N-fertilizer is lost (via NO_3^- leaching and gaseous-N emissions) from production systems, before the crop has a chance to absorb and assimilate it into plant-protein [46,105,106]. Annual economic loss from lost N-fertilizer is estimated at 90 US\$ billion [30,99]. If this trend continues, annual N-fertilizer application is expected to reach 300 Tg N by 2050; and the global N_2O emissions will reach 19 Tg y^{-1} by 2100 (from 10 Tg in 1990) [11,39,42,107]. There is an urgency to develop next-generation technologies to reduce N_2O emissions from agricultural systems as the IPCC set a target to cut global greenhouse gas emissions by 80% by 2100 [13]; EU, USA and China have committed to cut emissions by 30–40% (at the 1990 levels) by 2025.

Controlling soil nitrification is critical to reverse the present trend in declining NUE, and to improve N-retention and reduce N_2O emissions from agricultural systems. A paradigm shift is needed to move away from an inherently inefficient NO_3^- -centric nutrition and towards NH_4^+ -centric crop nutrition. Synthetic nitrification inhibitors are neither cost-effective nor functionally stable [30,60,89]. BNI-function, where plant-root systems deliver powerful BNIs at nitrifier-sites, should be genetically exploited as a plant trait for developing low-nitrifying, low- N_2O emitting next-generation N-efficient production systems.

Though, early observations of nitrification inhibition were mostly made on tropical grassland systems, BNI function seems not confined to plants either from humid- or sub-humid tropics as certain temperate forest ecosystems (such as pine forests) also suppress nitrification; selected temperate grasses likely also have BNI capacity in their root systems (recent unpublished reports and personal communications); also *Brassicaceae* members that are adapted to temperate climate have BNIs (isothiocyanates) in their root and shoot tissues and probably release these BNIs into the soil as well. In addition, some of the wheat wild relatives have high BNI-capacity, further suggesting that BNI function is widespread in both tropical and temperate plants.

High BNI-capacity root systems can be developed in both tropical- and temperate- crops and pastures using classical and modern breeding tools and approaches. Nitrification inhibitor producing plants such as *Brassicaceae* members can be incorporated in the soils (similar to green manures) and cropping and rotations can be developed with the primary objective of controlling soil nitrifier activity to improve NUE of production systems. High-BNI capacity *Brachiaria* pastures can be integrated with low-BNI capacity crops such as maize or upland rice in agro-pastoral systems which could fit well with the current move towards ecological-intensified agriculture. Low-nitrifying soil environment is an essential requirement for reducing N_2O emissions from agricultural systems and to limit nitrogen leakage into the larger environment. However, low-nitrifying soil environments must be complemented with restoration of key microbial communities to facilitate synchronization between SOM mineralization and crop-N demand [34]. Genetic and agronomic exploitation of BNI function in crops and pastures can facilitate moving towards low-nitrifying and low- N_2O emitting agricultural systems which can be an integral part of second Green Revolution.

Acknowledgements

Funding for BNI research at JIRCAS is mostly provided by MAFF (Japanese Ministry of Agriculture, Forest and Fisheries) and JSPS

research grant on wheat-BNI. The research on BNI at CIAT is supported by BMZ-GIZ, Germany; MADR, Colombia; MOFA and MAFF, Japan; and SIDA, Sweden.

References

- [1] R.A. Berner, GEOCARBSULF: a combined model for phanerozoic atmospheric O_2 and CO_2 , *Geochim. Cosmochim. Acta* 70 (2006) 5653–5664.
- [2] J.I. Prosser, Autotrophic nitrification in bacteria, *Adv. Microb. Physiol.* 30 (1989) 125–181.
- [3] S. Leninger, et al., Archaea predominate among ammonia-oxidizing prokaryotes in soils, *Nature* 442 (2006) 806–809.
- [4] J. Hahn, P.J. Crutzen, The role of fixed nitrogen in atmospheric phytochemistry, *Phil. Trans. Royal Soc. (Lond.) Ser. B* 296 (1982) 521–541.
- [5] C. Kroeze, Nitrous oxide and global warming, *Sci. Total Env.* 143 (1994) 193–209.
- [6] B.A. Hungate, J.S. Dukes, M.R. Shaw, Y. Luo, C.B. Field, Nitrogen and climate change, *Science* 302 (2003) 1512–1513.
- [7] J.M. Bremner, A.M. Blackmer, Nitrous oxide: emission from soils during nitrification of fertilizer nitrogen, *Science* 199 (1978) 295–296.
- [8] K.A. Smith, I.P. McTaggart, H. Tsuruta, Emissions of N_2O and NO associated with nitrogen fertilization in intensive agriculture, and the potential for mitigation, *Soil Use Manage.* 13 (1997) 296–304.
- [9] F.N. Tubiello, et al., The FAOSTAT database of greenhouse gas emissions from agriculture, *Env. Res. Lett.* 8 (2013), <http://dx.doi.org/10.1088/1748-9326/8/1/015009>.
- [10] J.N. Galloway, et al., Transformation of the nitrogen cycle: recent trends, questions, and potential solutions, *Science* 320 (2008) 889–892.
- [11] W.H. Schlesinger, On the fate of anthropogenic nitrogen, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 203–208.
- [12] L. Philippot, S. Hallin, Towards food, feed and energy crops mitigating climate change, *Trends Plant Sci.* 16 (2011) 476–480.
- [13] Intergovernmental Panel on Climate Change (IPCC), Fifth Assessment Synthesis Report, World Meteorological Organization/United Nations Environ, Prog., Paris, 2014.
- [14] C.F. Jordan, R.L. Todd, G. Escalante, Nitrogen conservation in a tropical rain forest, *Oecologia* 39 (1979) 123–128.
- [15] A.H. Magill, Long-term nitrogen additions and nitrogen saturation in two temperate forests, *Ecosystems* 3 (2000) 238–253.
- [16] I.T. Baldwin, R.K. Olson, W.A. Reinert, Protein binding phenolics and the inhibition of nitrification in subalpine balsam-fir soils, *Soil Biol. Biochem.* 15 (1983) 419–423.
- [17] P.R. Northup, Y. Zengshou, R.A. Dahlgren, K.A. Vogt, Polyphenol control of nitrogen release from pine litter, *Nature* 377 (1995) 227–229.
- [18] A. Smolander, S. Kanerva, B. Adamezyk, V. Kitunen, Nitrogen transformations in boreal forest soils – does composition of plant secondary compounds give any explanations, *Plant Soil* 350 (2012) 1–26.
- [19] P.M. Vitousek, P.A. Matson, Mechanisms of nitrogen retention in forest ecosystems: a field experiment, *Science* 225 (1984) 51–52.
- [20] A.B. Cooper, Suppression of nitrate formation with an exotic conifer plantation, *Plant Soil* 93 (1986) 383–394.
- [21] H. Stelzer, W.D. Bowman, Differential influence of plant species on soil nitrogen transformations within moist meadow alpine tundra, *Ecosystems* 1 (1988) 464–474.
- [22] C.S. White, The role of monoterpenes in soil nitrogen cycling processes in ponderosa pine; results from laboratory bioassays and field studies, *Biogeochemistry* 12 (1991) 43–68.
- [23] I. Ashton, W.A.E. Miller, W.D. Bowman, K.N. Suding, Niche complementarity due to plasticity in resource use: plant partitioning of chemical N forms, *Ecology* 91 (2010) 3252–3260.
- [24] E. Rice, S.K. Panchoy, Inhibition of nitrification by climax ecosystems, *Am. J. Bot.* 59 (1972) 1033–1040.
- [25] E. Rice, S.K. Panchoy, Inhibition of nitrification by climax ecosystems III. Inhibitors other than tannins, *Am. J. Bot.* 61 (1974) 1095–1103.
- [26] T. Nasholm, et al., Boreal forest plants take up organic nitrogen, *Nature* 392 (1998) 914–916.
- [27] L. Paavolainen, V. Kitunen, A. Smolander, Inhibition of nitrification in forest soil by monoterpenes, *Plant Soil* 205 (1998) 147–154.
- [28] K.G. Cassman, A.R. Dobermann, D.T. Walters, Agro-ecosystems, nitrogen-use efficiency and nitrogen management, *AMBIO* 31 (2002) 132–140.
- [29] R.J. Haynes, Mineral Nitrogen in the Plant–Soil System, Academic Press Inc., London, 1986, pp. 149–155.
- [30] G.V. Subbarao, et al., A paradigm shift towards low-nitrifying production systems: the role of biological nitrification inhibition (BNI), *Ann. Bot.* 112 (2013) 297–316.
- [31] D.L. Dinnes, et al., Nitrogen management strategies to reduce nitrate leaching in tile drained Mid-Western soils, *Agron. J.* 94 (2002) 153–171.
- [32] D. Tilman, K.G. Cassman, P.A. Matson, R. Naylor, S. Polasky, Agricultural sustainability and intensive production practices, *Nature* 418 (2002) 671–677.
- [33] A.E. Russell, C.A. Cambardella, D.A. Laird, D.B. Jaynes, D.W. Meek, Nitrogen fertilizer effects on soil carbon balances in Midwestern US agricultural systems, *Ecol. Appl.* 19 (2009) 1102–1113.

- [34] N. Perveen, et al., Priming effect and microbial diversity in ecosystem functioning and response to global change: a modeling approach using the SYMPHONY model, *Global Change Biol.* 20 (2014) 1174–1190.
- [35] A. Fortuna, R.R. Harwood, G.P. Robertson, J.W. Fisk, E.A. Paul, Seasonal changes in nitrification potential associated with application of N fertilizer and compost in maize systems of southwest Michigan, *Agric. Ecosyst. Environ.* 97 (2003) 285–293.
- [36] H. Chu, T. Fujii, S. Morimoto, X. Lin, K. Yagi, J. Hu, J. Jhang, Community structure of ammonia-oxidizing bacteria under long-term application of mineral fertilizer and organic manure in a sandy loam soil, *Appl. Environ. Microbiol.* 73 (2007) 485–491.
- [37] J.M. Norton, J.M. Stark, Regulation and measurement of nitrification in terrestrial systems, *Methods Enzymol.* 486 (2011) 343–368.
- [38] P.M. Vitousek, et al., Human alteration of the global nitrogen cycle: sources and consequences, *Ecol. Appl.* 7 (1997) 737–750.
- [39] D. Tilman, et al., Forecasting agriculturally driven global environmental change, *Science* 292 (2001) 281–284.
- [40] G.V. Subbarao, et al., Nitrogen management in grasslands in forage-based production systems – Role of biological nitrification inhibition (BNI), *Trop. Grasslands* 1 (2013) 168–174.
- [41] E. Kahril, et al., Greenhouse gas emissions from nitrogen use in China, *Environ. Sci. Policy* 13 (2010) 688–694.
- [42] J. Rockstrom, et al., A safe operating space for humanity, *Nature* 461 (2009) 472–475.
- [43] R.J. Haynes, K.M. Goh, Ammonium and nitrate nutrition of plants, *Biol. Rev.* 53 (1978) 465–510.
- [44] S. Boudsocq, J.C. Lata, J. Mathieu, L. Abbadié, S. Barot, Modelling approach to analyse the effects of nitrification inhibition on primary production, *Funct. Ecol.* 23 (2009) 220–230.
- [45] H. Marschner, Mineral Nutrition of Higher Plants, 2nd Ed., Academic Press, London, 1995.
- [46] W.R. Raun, G.V. Johnson, Improving nitrogen use efficiency for cereal production, *Agron. J.* 91 (1999) 357–363.
- [47] A.D.M. Glass, Nitrogen use efficiency of crop plants: physiological constraints upon nitrogen absorption, *Crit. Rev. Plant Sci.* 22 (2003) 453–470.
- [48] A.C. Finzi, et al., Increases in nitrogen uptake rather than nitrogen-use efficiency support higher rates of temperate forest productivity under elevated CO₂, *Proc. Natl. Acad. Sci. U. S. A.* 104 (2007) 14014–14019.
- [49] G.V. Subbarao, et al., Evidence for biological nitrification inhibition in *Brachiaria* pastures, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 17302–17307.
- [50] J.C. Lata, J. Durand, R. Lensi, L. Abbadié, Stable coexistence of contrasted nitrification statuses in a wet tropical savanna ecosystem, *Funct. Ecol.* 13 (1999) 762–768.
- [51] J.C. Lata, Interactions between Microbial Processes, Nutrient cycle and Grass cover Functioning: Study of Soil Nitrification under the Gramineae *Hyparrhenia diplandra* in a Wet Tropical Savanna of Ivory Coast, Univ. Paris, France, 1999 (Ph.D Thesis).
- [52] J.C. Lata, et al., Grass populations control nitrification in savanna soils, *Funct. Ecol.* 18 (2004) 605–611.
- [53] S. Boudsocq, et al., Plant preference for ammonium versus nitrate; a neglected determinant of ecosystem functioning? *Am. Nat.* 180 (2012) 60–69.
- [54] T. Iizumi, M. Mizumoto, K. Nakamura, A bioluminescence assay using *Nitrosomonas europaea* for rapid and sensitive detection of nitrification inhibitors, *Appl. Environ. Microbiol.* 64 (1998) 3656–3662.
- [55] G.V. Subbarao, et al., A bioluminescence assay to detect nitrification inhibitors released from plant roots: a case study with *Brachiaria humidicola*, *Plant Soil* 288 (2006) 101–112.
- [56] S.C. Hart, J.M. Stark, E.A. Davidson, M.K. Firestone, Nitrogen mineralization, immobilization, and nitrification, in: R.W. Hattori, J.S. Angle, B.S. Bottomley (Eds.), *Methods of Soil Analysis, Part 2. Microbiological and Biochemical Properties*, Soil Sci. Soc. Am., Madison, 1994, pp. 985–1018.
- [57] N.A.C. Smits, R. Bobbink, H.J. Laanbroek, A.J. Paalman, M.M. Hefting, Repression of potential nitrification activities by matgrass sward species, *Plant Soil* 337 (2010) 435–445.
- [58] G.V. Subbarao, et al., Biological nitrification inhibition (BNI) – is it a widespread phenomenon? *Plant Soil* 294 (2007) 5–18.
- [59] G.V. Subbarao, H.Y. Wang, O. Ito, K. Nakahara, W.L. Berry, NH₄⁺ triggers the synthesis and release of biological nitrification inhibition compounds in *Brachiaria humidicola* roots, *Plant Soil* 290 (2007) 245–257.
- [60] G.V. Subbarao, et al., Scope and strategies for regulation of nitrification in agricultural systems – challenges and opportunities, *Critic. Rev. Plant Sci.* 25 (2006) 303–335.
- [61] T.E. Crews, The presence of nitrogen fixing legumes in terrestrial communities: evolutionary vs ecological considerations, *Biogeochemistry* 46 (1999) 233–246.
- [62] G.V. Subbarao, et al., Free fatty acids from the pasture grass *Brachiaria humidicola* and one of their methyl esters as indicators of nitrification, *Plant Soil* 313 (2008) 89–99.
- [63] H.A.K.M. Zakir, et al., Detection, isolation and characterization of a root-exuded compound, methyl 3-(4-hydroxyphenyl)propionate, responsible for biological nitrification inhibition by sorghum (*Sorghum bicolor*), *New Phytol.* 180 (2008) 442–451.
- [64] S. Gopalakrishnan, et al., Nitrification inhibitors from the root tissues of *Brachiaria humidicola*, a tropical grass, *J. Agric. Food Chem.* 55 (2007) 1385–1388.
- [65] G.V. Subbarao, et al., Biological nitrification inhibition (BNI) activity in sorghum and its characterization, *Plant Soil* 366 (2013) 243–259.
- [66] T. Tesfamariam, et al., Biological nitrification inhibition in sorghum: the role of sorgoleone production, *Plant Soil* 379 (2014) 325–335.
- [67] C.S. White, Nitrification inhibition by monoterpenoids: theoretical mode of action based on molecular structures, *Ecology* 69 (1988) 1631–1633.
- [68] G.D. Bending, S.D. Lincoln, Inhibition of soil nitrifying bacteria communities and their activities by glucosinolate hydrolysis products, *Soil Biol. Biochem.* 32 (2000) 1261–1269.
- [69] P.B. Brown, M.J. Morra, Brassicaceae tissues as inhibitors of nitrification in soil, *J. Agric. Food Chem.* 57 (2009) 7706–7711.
- [70] A.J. Snyder, J.L. Johnson-Maynard, M.J. Morra, Nitrogen mineralization in soil incubated with 15N-labeled Brassicaceae seed meals, *Appl. Soil Ecol.* 46 (2010) 73–80.
- [71] X. Yang, B.E. Scheffler, L.A. Weston, SOR1, a gene associated with bioherbicide production in sorghum root hairs, *J. Exp. Bot.* 55 (2004) 2251–2259.
- [72] S.R. Baerson, A.M. Rimando, Z. Pan, Probing allelochemical biosynthesis in sorghum root hairs, *Plant Signal. Behav.* 3 (2008) 667–670.
- [73] A.H. de Boer, I.J. de Vries-van Leeuwen, Fusicocanones: diterpenes with surprising biological functions, *Trends Plant Sci.* 17 (2012) 360–368.
- [74] G.S. Muromtsev, et al., Occurrence of fusicocanones in plants and fungi, *J. Plant Growth Reg.* 13 (1994) 39–49.
- [75] T. Toyomasu, et al., Fusicocins are biosynthesized by an unusual chimera synthase in fungi, *Proc. Natl. Acad. Sci. U. S. A.* 104 (2007) 3084–3088.
- [76] A. Osbourn, Secondary metabolic gene clusters: evolutionary toolkits for chemical innovation, *Trends Genet.* 26 (2010) 449–457.
- [77] A. Osbourn, Gene clusters for secondary metabolic pathways: an emerging theme in plant biology, *Plant Physiol.* 154 (2010) 532–535.
- [78] H.Y. Chu, E. Wegel, A. Osbourn, From hormones to secondary metabolism: the emergence of secondary metabolic gene clusters in plants, *Plant J.* 66 (2011) 66–79.
- [79] A. Okada, et al., *OsTGAP1*, a *bZIP* transcription factor coordinately regulates the inductive production of diterpenoid phytoalexins in rice, *J. Biol. Chem.* 284 (2009) 26510–26518.
- [80] K. Okada, The biosynthesis of isoprenoids and the mechanisms regulating it in plants, *Biosci. Biotechnol. Biochem.* 75 (2011) 1219–1225.
- [81] T. Shimizu, et al., The potential bio-production of the pharmaceutical agent sakuranetin, a flavonoid phytoalexin in rice, *Bioengineered* 4 (2012) 352–357.
- [82] F.E. Dayan, et al., Sorgoleone, *Phytochemistry* 71 (2010) 1032–1039.
- [83] F.E. Dayan, J. Howell, J.D. Weidenhamer, Dynamic root exudation of sorgoleone and its mechanism of action, *J. Exp. Bot.* 60 (2009) 2107–2117.
- [84] G.V. Subbarao, et al., Biological nitrification inhibition (BNI) – is there potential for genetic interventions in the Triticeae, *Breed. Sci.* 59 (2009) 529–545.
- [85] G.W. McCarty, Modes of action of nitrification inhibitors, *Biol. Fertil. Soils* 29 (1999) 1–9.
- [86] M.J. Fisher, et al., Carbon storage by introduced deep-rooted grasses in the South American savannas, *Nature* 371 (1994) 236–238.
- [87] G.V. Subbarao, et al., Can biological nitrification inhibition (BNI) genes from perennial *Leymus racemosus* (Triticeae) combat nitrification in wheat farming, *Plant Soil* 299 (2007) 55–64.
- [88] Y. Zhu, H. Zeng, Q. Shen, T. Ishikawa, G.V. Subbarao, Interplay among NH₄⁺ uptake, rhizosphere pH and plasma membrane H⁺-ATPase determine the release of BNIs in sorghum roots – possible mechanisms and underlying hypothesis, *Plant Soil* 358 (2012) 131–141.
- [89] G.V. Subbarao, et al., Biological nitrification inhibition – a novel strategy to regulate nitrification in agricultural systems, *Adv. Agron.* 114 (2012) 249–302.
- [90] S. Gopalakrishnan, et al., Biological nitrification inhibition (BNI) by *Brachiaria humidicola* roots varies with soil type and inhibits nitrifying bacteria but not other soil microorganisms, *Soil Sci. Plant Nutr.* 55 (2009) 725–733.
- [91] L.M. Zahn, A boost from wild wheat, *Science* 318 (2007) 171.
- [92] A. Mujeeb-Kazi, V. Rosas, S. Roldan, Conservation of the genetic variation of *Triticum tauschii* (Coss.) Schmalh. Lslat. \times *T. tauschii*; $2n = 6x = 42$, AABDD) and its potential utilization for wheat improvement, *Genet. Res. Crop Evol.* 43 (1996) 129–134.
- [93] A. Mujeeb-Kazi, et al., Current status of D-genome based synthetic hexaploid wheats, and the characterization of an elite subset, *Ann. Wheat Newsllett.* 46 (2000) 76–79.
- [94] T.S. Walker, H.P. Bais, E. Grotewold, J.M. Vivanco, Root exudation and rhizosphere biology, *Plant Physiol.* 132 (2003) 44–51.
- [95] D.E. Moreta, et al., Biological nitrification inhibition (BNI) in *Brachiaria* pastures: a novel strategy to improve eco-efficiency of crop-livestock systems and to mitigate climate change, *Trop. Grasslands* 2 (2014) 88–91.
- [96] A. Weigelt, R. Bol, R.D. Bardgett, Preferential uptake of soil nitrogen forms by grassland species, *Oecology* 142 (2005) 627–635.
- [97] M.A. Sutton, et al., Too much of a good thing, *Nature* 472 (2011) 159–161.
- [98] FAO, FAOSTAT Database – Agriculture Production, Food and Agriculture Organization of the United Nations, Rome, 2011.
- [99] R.L. Mulvaney, S.A. Khan, T.R. Ellsworth, Synthetic nitrogen fertilizers deplete soil nitrogen: a global dilemma for sustainable cereal production, *J. Environ. Qual.* 38 (2009) 2295–2314.
- [100] E.T. Elliott, Aggregate structure and carbon, nitrogen and phosphorus in native and cultivated soils, *Soil Sci. Soc. Am. J.* 50 (1986) 627–633.
- [101] S.M. Ross, Organic matter in tropical soils: current conditions, concerns and prospects for conservation, *Prog. Phys. Geo.* 17 (1993) 265–305.
- [102] I. Celik, Land-use effects on organic matter and physical properties of soil in a southern Mediterranean highland of Turkey, *Soil Till. Res.* 83 (2005) 270–277.

- [103] P.A. Matson, A.R. McDowell Townsend, P.M. Vitousek, The globalization of N deposition: ecosystem consequences in tropical environments, *Biogeochemistry* 46 (1999) 67–83.
- [104] G.V. Subbarao, I.M. Rao, K. Nakahara, K.L. Sahrawat, Y. Ando, T. Kawashima, Potential for biological nitrification inhibition to reduce nitrification and N₂O emissions in pasture crop–livestock systems, *Animal* 7 (s2) (2013) 322–332.
- [105] W.T. Peterjohn, W.H. Schlesinger, Nitrogen loss from deserts in the South Western United States, *Biogeochemistry* 10 (1990) 67–79.
- [106] P.M. Vitousek, R.W. Howarth, Nitrogen limitation on land and in the sea: how can it occur, *Biogeochemistry* 13 (1991) 87–115.
- [107] R.E. Turner, N.N. Rabalais, D. Justic, Gulf of Mexico hypoxia: alternate states and a legacy, *Environ. Sci. Technol.* 42 (2008) 2323–2327.